VOL. 630, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY FEBRUARY 5, 1993

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Retention on non-polar adsorbents in liquid–solid chromatography. Effect of grafted alkyl chains
by G. Fóti, M. L. Belvito, A. Alvarez-Zepeda and E. sz. Kováts (Lausanne, Switzerland) (Received October 26th, 1992) 1

Modeling of the adsorption behavior and the chromatographic band profiles of enantiomers. Behavior of methyl mandelate on immobilized cellulose
by F. Charton, S. C. Jacobson and G. Guiochon (Knoxville, TN and Oak Ridge, TN, USA) (Received October 27th, 1992) 21

Characterization of non-linear adsorption properties of dextran-based polyelectrolyte displacers in ion-exchange systems
by S. D. Graham, G. Jayaraman and S. M. Cramer (Troy, NY, USA) (Received October 1st, 1992) 37

Ion-exchange displacement chromatography of proteins. Dextran-based polyelectrolytes as high affinity displacers
by G. Jayaraman, S. D. Graham and S. M. Cramer (Troy, NY, USA) (Received October 1st, 1992) 53

Improved computer algorithm for characterizing skewed chromatographic band broadening. II. Results and comparisons
by W. W. Yau, S. W. Rementer, J. M. Boyajian, J. J. DeStefano, J. F. Graff, K. B. Lim and J. J. Kirkland (Wilmington, DE, USA) (Received October 14th, 1992) 69

Functionalized polymer particles for chiral separation
by T. Hargitai, P. Reinholdsson and B. Törnell (Lund, Sweden) and R. Isaksson (Uppsala, Sweden) (Received October 21st, 1992) 79

New alumina-based stationary phases for high-performance liquid chromatography. Synthesis by olefin hydrosilation on a silicon hydride-modified alumina intermediate
by J. J. Pesek, J. E. Sandoval and M. Su (San Jose, CA, USA) (Received October 16th, 1992) 95

Examination of some reversed-phase high-performance liquid chromatography systems for the determination of lipophilicity
by D. P. Nowotnik, T. Feld and A. D. Nunn (New Brunswick, NJ, USA) (Received September 29th, 1992) 105

Evaluation of silanol-deactivated silica-based reversed phases for liquid chromatography of erythromycin
by J. Paesen, P. Claeys, E. Roets and J. Hoogmartens (Leuven, Belgium) (Received October 12th, 1992) 117

Selective complex formation of saccharides with europium(III) and iron(III) ions at alkaline pH studied by ligand-exchange chromatography
by M. Stefansson (Uppsala, Sweden) (Received October 20th, 1992) 123

Separation of glucooligosaccharides and polysaccharide hydrolysates by gradient elution hydrophilic interaction chromatography with pulsed amperometric detection
by A. S. Feste and I. Khan (Houston, TX, USA) (Received October 23rd, 1992) 129

Determination of hyaluronic acid by high-performance liquid chromatography of the oligosaccharides derived therefrom as 1-(4-methoxy)phenyl-3-methyl-5-pyrazolone derivatives
by K. Kakehi, M. Ueda, S. Suzuki and S. Honda (Osaka, Japan) (Received September 23rd, 1992) 141

Separation and isolation of trace impurities in L-tryptophan by high-performance liquid chromatography
by M. W. Trucksess (Washington, DC, USA) (Received October 10th, 1992) 147

Sequence-specific DNA affinity chromatography: application of a group-specific adsorbent for the isolation of restriction endonucleases
by C. Pozidis, G. Vlatakis and V. Bouriotis (Heraklion, Greece) (Received October 9th, 1992) 151

Quantitative analysis of tylosin by column liquid chromatography
by E. Roets, P. Beirinckx, I. Quintens and J. Hoogmartens (Leuven, Belgium) (Received September 29th, 1992) . . . 159

Analysis of the explosive 2,4,6-trinitrophenylmethylnitramine (tetryl) in bush bean plants
by S. D. Harvey, R. J. Fellows, D. A. Cataldo and R. M. Bean (Richland, WA, USA) (Received September 29th, 1992) 167

(Continued overleaf)

Contents (continued)

Liquid–gel partitioning using Lipidex in the determination of polychlorinated biphenyls in cod liver oil
by C. Weistrand and K. Norén (Stockholm, Sweden) (Received October 19th, 1992) 179

High-performance liquid chromatographic determination of methanesulphinic acid as a method for the determination of hydroxyl radicals
by S. Fukui, Y. Hanasaki and S. Ogawa (Kyoto, Japan) (Received September 24th, 1992) 187

Ion chromatography of nitrite and carbonate in inorganic matrices on an octadecyl–poly(vinyl alcohol) gel column using acidic eluents
by S. Rokushika, F. M. Yamamoto and K. Kihara (Kyoto, Japan) (Received October 13th, 1992) 195

Column Liquid Chromatography–Gas Chromatography

Use of an open-tubular trapping column as phase-switching interface in on-line coupled reversed-phase liquid chromatography–capillary gas chromatography
by H. G. J. Mol, J. Staniewski, H.-G. Janssen and C. A. Cramers (Eindhoven, Netherlands) and R. T. Ghijsen and U. A. Th. Brinkman (Amsterdam, Netherlands) (Received October 12th, 1992) 201

Comparative study of the determination of triacylglycerol in vegetable oils using chromatographic techniques
by A. A. Carelli and A. Cert (Seville, Spain) (Received October 20th, 1992) 213

Gas Chromatography

Structure–gas chromatographic retention time models of tetra-*n*-alkylsilanes and tetra-*n*-alkylgermanes using topological indexes
by E. J. Kupchik (Jamaica, NY, USA) (Received October 27th, 1992) 223

Effect of variations in gas chromatographic conditions on the linear retention indices of selected chemical warfare agents
by M. Kokko (Helsinki, Finland) (Received September 23rd, 1992) 231

Relationship between Kováts retention indices and molecular connectivity indices of tetralones, coumarins and structurally related compounds
by A. C. Arruda, V. E. F. Heinzen and R. A. Yunes (Florianópolis, Brazil) (Received August 18th, 1992) 251

Ambient temperature gas purifier suitable for the trace analysis of carbon monoxide and hydrogen and the preparation of low-level carbon monoxide calibration standards in the field
by B. E. Foulger and P. G. Simmonds (Poole, UK) (Received August 7th, 1992) 257

Evaluation of Carboxen carbon molecular sieves for trapping replacement chlorofluorocarbons
by S. J. O'Doherty, P. G. Simmonds and G. Nickless (Bristol, UK) and W. R. Betz (Bellefonte, PA, USA) (Received October 29th, 1992) 265

Application of dithiocarbamate resin–metal complexes as stationary phases in gas chromatography
by C.-F. Yeh, S.-D. Chyueh, W.-S. Chen, J.-D. Fang and C.-Y. Liu (Taipei, Taiwan) (Received October 20th, 1992) . 275

Gas chromatographic analysis of high-molecular-mass polycyclic aromatic hydrocarbons. II. Polycyclic aromatic hydrocarbons with relative molecular masses exceeding 328
by A. Bemgård, A. Colmsjö and B.-O. Lundmark (Solna, Sweden) (Received October 15th, 1992) 287

Simultaneous determination of planar chlorobiphenyls and polychlorinated dibenzo-*p*-dioxins and -furans in Dutch milk using isotope dilution and gas chromatography–high-resolution mass spectrometry
by J. A. van Rhijn, W. A. Traag, P. F. van de Spreng and L. G. M. Th. Tuinstra (Wageningen, Netherlands) (Received August 19th, 1992) 297

Supercritical Fluid Chromatography

Injection of large volumes of aqueous solutions in capillary supercritical fluid chromatography and sample preconcentration by multiple injections
by S. Bouissel, F. Erni and R. Link (Basle, Switzerland) (Received September 28th, 1992) 307

Electrophoresis

pH gradient simulator for electrophoretic techniques in a Windows environment
by E. Giaffreda, C. Tonani and P. G. Righetti (Milan, Italy) (Received October 19th, 1992) 313

Protein and peptide mobility in capillary zone electrophoresis. A comparison of existing models and further analysis
by V. J. Hilser, Jr. and G. D. Worosila (Suffern, NY, USA) and S. E. Rudnick (Riverdale, NY, USA) (Received November 2nd, 1992) 329

Capillary electrophoretic protein separations in polyacrylamide-coated silica capillaries and buffers containing ionic surfactants
by M. A. Strege and A. L. Lagu (Indianapolis, IL, USA) (Received October 30th, 1992) 337

Determination of the number and relative molecular mass of subunits in an oligomeric protein by two-dimensional electrophoresis. Application to the subunit structure analysis of rat liver amidophosphoribosyltransferase
by T. Yamaoka and K. Yamashita (Ibaraki, Japan) and M. Itakura (Tokushima City, Japan) (Received October 6th, 1992) 345

Separation of drug stereoisomers by capillary electrophoresis with cyclodextrins
by T. E. Peterson (Fort Worth, TX, USA) (Received October 21st, 1992) 353

Separation and simultaneous determination of the active ingredients in theophylline tablets by micellar electrokinetic capillary chromatography
by Q.-x. Dang and L.-x. Yan (Xian, China) and Z.-p. Sun and D.-k. Ling (Beijing, China) (Received October 20th, 1992) 363

Determination of alkylphosphonic acids by capillary zone electrophoresis using indirect UV detection
by G. A. Pianetti, M. Taverna, A. Baillet, G. Mahuzier and D. Baylocq-Ferrier (Chatenay-Malabry, France) (Received October 23rd, 1992) 371

Metal ion capillary zone electrophoresis with direct UV detection: determination of transition metals using an 8-hydroxyquinoline-5-sulphonic acid chelating system
by A. R. Timerbaev, W. Buchberger, O. P. Semenova and G. K. Bonn (Linz, Austria) (Received August 11th, 1992) . 379

SHORT COMMUNICATIONS

Column Liquid Chromatography

Glycosaminoglycans and proteins: different behaviours in high-performance size-exclusion chromatography
by N. Volpi and L. Bolognani (Modena, Italy) (Received October 28th, 1992) 390

Adsorption losses during extraction and derivatization efficiency by benzoylation of plant putrescine for high-performance liquid chromatographic analysis
by M. Z. Hauschild (Lyngby, Denmark) (Received November 2nd, 1992) 397

High-performance liquid chromatographic determination of cardenolides in *Digitalis* leaves after solid-phase extraction
by H. Wiegrebe and M. Wichtl (Marburg/Lahn, Germany) (Received October 27th, 1992) 402

Determination of acidic saponins in crude drugs by high-performance liquid chromatography on octadecylsilyl porous glass
by H. Kanazawa, Y. Nagata, Y. Matsushima, M. Tomoda and N. Takai (Tokyo, Japan) (Received November 3rd, 1992) 408

Determination of Lovastatin (mevinolin) and mevinolinic acid in fermentation liquids
by R. Kysilka and V. Křen (Prague, Czechoslovakia) (Received October 23rd, 1992) 415

Simultaneous high-performance liquid chromatographic determination of alterosolanol A, B, C, D, E and F
by N. Okamura, A. Yagi, H. Haraguchi and K. Hashimoto (Hiroshima, Japan) (Received November 10th, 1992) . . 418

Isolation and determination of alizarin in cell cultures of *Rubia tinctorum* and emodin in *Dermocybe sanguinea* using solid-phase extraction and high-performance liquid chromatography
by Z. A. Tóth (Budapest, Hungary) and O. Raatikainen, T. Naaranlahti and S. Auriola (Kuopio, Finland) (Received October 28th, 1992) 423

Gas Chromatography

Improved procedure for the derivatization and gas chromatographic determination of hydroxycarboxylic acids treated with chloroformates
by P. Hušek (Prague, Czechoslovakia) (Received October 9th, 1992) 429

Preparation of methyl esters of fatty acids with trimethylsulphonium hydroxide —an appraisal
by A. H. El-Hamdy and W. W. Christie (Ayr, UK) (Received November 12th, 1992) 438

Electrophoresis

Capillary electrophoretic separation of recombinant granulocyte-colony-stimulating factor glycoforms
by E. Watson and F. Yao (Thousand Oaks, CA, USA) (Received November 23rd, 1992) 442

BOOK REVIEW

Advances in lipid methodology 1 (edited by W. W. Christie), by A. Kuksis (Toronto, Canada) 447

AUTHOR INDEX 449